

Non-abelian Berry's phase and Chern numbers in higher spin pairing condensates

Chyh-Hong Chern,¹ Han-Dong Chen,² Congjun Wu,¹ Jiang-Ping Hu,³ and Shou-Cheng Zhang¹

¹*Department of Physics, McCullough Building, Stanford University, Stanford CA 94305-4045*

²*Department of Applied Physics, McCullough Building, Stanford University, Stanford CA 94305-4045*

³*Department of Physics and Astronomy, University of California, Los Angeles, CA 90095-1547*

We show that the non-Abelian Berry phase emerges naturally in the s-wave and spin quintet pairing channel of spin-3/2 fermions. The topological structure of this pairing condensate is characterized by the second Chern number. This topological structure can be realized in ultra-cold atomic systems and in solid state systems with at least two Kramers doublets.

PACS numbers: 74.90.+n, 71.10.Fd, 03.75.Nt, 02.40.Re

I. INTRODUCTION

Topological gauge structure and Berry's phase¹ play an increasingly important role in condensed matter physics. The quantized Hall conductance can be deeply understood in terms of the first Chern class^{2,3}. The fractional quantum Hall effect can be fundamentally described by the $U(1)$ topological Chern-Simons gauge theory⁴. The effective action for ferromagnets⁵ and one dimensional antiferromagnets^{6,7} contain Berry phase terms which fundamentally determine the low energy dynamics. More recently, Berry's phase associated with the BCS quasi-particles in pairing condensates has also been studied extensively.^{8,9,10}

However, most of the well known applications of the geometrical phase in condensed matter physics involves Berry's¹ original abelian $U(1)$ phase factor. More recently, the non-abelian $SU(2)$ ^{11,12,13} Berry's phase (or holonomy, to be precise) has been systematically investigated in the context of condensed matter systems. Demler and Zhang¹⁴ investigated the quasi-particle wave functions in the unified $SO(5)$ theory of antiferromagnetism and superconductivity, and found that the SDW and the BCS quasi-particle states accumulate an $SU(2)$ Berry's phase (or holonomy) when the order parameter returns to itself after an adiabatic circuit. Zhang and Hu¹⁵ found a higher dimensional generalization of the quantum Hall effect based on a topologically non-trivial $SU(2)$ background gauge field. Rather surprisingly, the non-abelian $SU(2)$ holonomy also found its deep application in the technologically relevant field of quantum spintronics^{16,17}. All these condensed matter applications are underpinned by a common mathematical framework, which naturally generalizes the concept of Berry's $U(1)$ phase factor. This new class of applications is topologically characterized by the second Chern class and the second Hopf map, and applies to fermionic systems with time reversal invariance.

In this paper, we investigate the non-trivial topological structures associated with the higher spin condensates. We first review the momentum space gauge structure of the spin one condensate, namely the A phase of 3He . As it has been pointed out⁸, the momentum space gauge structure of the pairing condensate is given by that of

the t'Hooft-Polyakov monopole. We then investigate the new system of spin 2 (quintet) pairing condensate of the underlying spin 3/2 fermions. The most general Hubbard model of spin 3/2 fermions has recently been introduced and investigated extensively by Wu et al.¹⁸, who found that the model always has a generic $SO(5)$ symmetry in the spin sector. Building on this work, we show here that the fermionic quasi-particles of the quintet pairing condensate can be described by the second Hopf map. Similar to the SDW+BCS system investigated by Demler and Zhang¹⁴, the quasi-particles of the quintet pairing condensate also accumulate an $SU(2)$ holonomy. The quintet pairing condensate can be experimentally realized in a number of systems. Cold atoms with spin 3/2 in the continuum or on the optical lattice can be accurately described by the model of local contact interactions¹⁸ U_0 and U_2 . These interaction parameters can be experimentally tunes over a wide range, including the range for stable quintet condensates. Effective spin 3/2 fermions can also be realized in solid state systems with at least two Kramers doublets, for example in bands formed by $P_{3/2}$ orbitals.

In the rest of this paper, we shall use spin-1/2 system to be short for the spin-1/2 superfluid 3He -A and spin-3/2 system for the s-wave spin-3/2 superconductor in the quintet channel. The repeated indices are summed assumably throughout this paper.

II. SUPERFLUID 3He -A

A. Goldstone manifold and the 1st Hopf map

The general form of the equilibrium order parameter in the 3He -A phase can be written as⁸

$$\langle \Delta(k)_{ai} \rangle = \Delta_k \hat{d}_a \left(\hat{e}_i^{(1)} + i \hat{e}_i^{(2)} \right), \quad (1)$$

where the spin index (a) and orbital index (i) run from 1 to 3. Δ_k is a complex number that contains the information of the magnitude and the $U(1)$ phase. $\hat{d}$ is the normal vector of the plane to which the spin direction is restricted. The orthogonal vectors $\hat{e}^{(1)}$, $\hat{e}^{(2)}$ and $\hat{l} = \hat{e}^{(1)} \times \hat{e}^{(2)}$ form a local physical coordinate frame.

The Goldstone manifold of the order parameter is given by⁸

$$\begin{aligned} R_a &= G/H = \frac{U(1) \times SO(3)^{(L)} \times SO(3)^{(S)}}{SO(2)^{(S)} \times U(1)^{\text{combined}} \times Z_2^{\text{combined}}} \\ &= S^2 \times SO(3)_{\text{relative}}/Z_2. \end{aligned} \quad (2)$$

Here $SO(3)_{\text{relative}}$ denotes such rotations about the axis $\hat{l}$ that lead to new degenerate states which are relative to gauge transformations⁸. The $U(1)^{\text{combined}}$ comes from the fact that the A-phase state is invariant under combined transformation⁸ of the gauge transformation with the parameter ϕ from the $U(1)$ group and the orbital rotation of $\hat{e}^{(1)}$, $\hat{e}^{(2)}$ and $\hat{l}$ about axis $\hat{l}$ by the same angle ϕ . The Z_2^{combined} denotes the combined operation that $\hat{d} \rightarrow -\hat{d}$, $\Delta_k \rightarrow -\Delta_k$. This combined discrete symmetry leads to the existence of half-quantum vortices⁸. Around a half-quantum vortex, the vector field $\hat{d}$ is continuously rotated into $-\hat{d}$ and the $U(1)$ phase of Δ_k continuously evolves from 0 to π when the order parameter returns to itself after an adiabatic circuit.

If we fix the local orthogonal frame in an arbitrary direction and adiabatically move the quasi-particle around a line defect of half-quantum vortex, the trajectory of the order parameter is a closed loop on the S^2/Z_2 space. On the other hand, the degrees of freedom of the quasi-particle (a 2-dimensional spinor) forms a 3-dimensional sphere S^3 and the trajectory of the quasi-particle on S^3 is not closed. This adiabatic evolution defines the following map

$$S^3 \longrightarrow S^2/Z_2. \quad (3)$$

In the topological terminology, Eq.(3) is determined by the third homotopic group denoted by $\pi_3(S^2/Z_2)$. Due to a theorem in the Homotopy theory¹⁹

$$\pi_k(S^n/Z_2) = \pi_k(S^n), \text{ for } k \geq 2. \quad (4)$$

Eq.(3) is homotopically equivalent to the first Hopf map $S^3 \rightarrow S^2$ i.e., $U(1)$ Berry phase in the FQHE and other nano-structure in the semiconductors^{20,21}.

B. Berry connection, 1st Chern number, t'Hooft-Polyakov monopole and Dirac monopole

If we define the spinor as

$$\Psi_k^\dagger = \left(c_{k, \frac{1}{2}}^\dagger, c_{k, -\frac{1}{2}}^\dagger, c_{-k, \frac{1}{2}}, c_{-k, -\frac{1}{2}} \right), \quad (5)$$

the mean field Hamiltonian for $^3\text{He-A}$ is given by

$$\mathcal{H} = \sum_k \Psi_k^\dagger H_k \Psi_k \quad (6)$$

with

$$H_k = \begin{pmatrix} \epsilon_k \sigma^0 & \Delta_k \\ \Delta_k^\dagger & -\epsilon_k \sigma^0 \end{pmatrix}, \quad (7)$$

where $\Delta_k = -\Delta_k \hat{d}_a \sigma^a R$ and $R = -i\sigma^2$. ϵ_k is the kinetic energy on the lattices referenced from the Fermi surface and the summation of momentum k is over half of the Brillouin zone to avoid the double counting. Here, σ^0 is the 2×2 identity matrix and $\sigma^{1,2,3}$ are Pauli matrices.

The Berry phase connection (BPC) is defined by the differential change of states projecting to themselves. In this paper, it will be illustrated by using the state with a positive eigenvalue. BPC obtained from the state with a negative eigenvalue is simply the complex conjugate to the one with the eigenvalue of a different sign. The BPC and its field strength can be obtained respectively.

$$A_a = -iA_a^c \frac{\sigma_c}{2}, \quad A_a^c = \epsilon_{abc} \frac{\hat{d}_b}{d}. \quad (8)$$

and

$$\begin{aligned} F_{bc}^a &= \partial_b A_c^a - \partial_c A_b^a + \epsilon_{ade} A_b^d A_c^e \\ &= -\frac{1}{d^2} \epsilon_{bce} \hat{d}_e \hat{d}_a, \end{aligned} \quad (9)$$

The gauge invariant magnetic field can be defined as

$$B^a = \frac{1}{2} \epsilon_{abc} F_{bc}^e d_e = -\frac{\hat{d}_a}{d^2}. \quad (10)$$

This is a $U(1)$ magnetic-monopole like field in the d -space. It emerges when there are line defects in the $\hat{d}$ field, e.g., half-quantum vortices in the superfluid $He-3A$ phase. If we transport the spin-1/2 fermion adiabatically around the vortex, the electronic wavefunction gains the phase accumulated due to the $\hat{d}$ field, as we discussed previously. Moreover, the first Chern number can be computed easily.

$$\mathcal{C}_1 = \frac{1}{4\pi} \oint \vec{B} \cdot \vec{d}S = -1. \quad (11)$$

This is the famous t'Hooft-Polyakov monopole (TPM)^{8,22}. Different from the Dirac monopole, the gauge field of TPM is non-abelian and finite everywhere over S^2/Z_2 while the Dirac magnetic monopole is abelian and has a singularity string. There is a deep and direct relation between them, which can be achieved by a singular gauge transformation^{23,24,25}.

The present $SO(3)$ Berry phase defines a $SO(3)$ gauge theory on S^2/Z_2 . Using the covariant derivative $D_a = \partial_a + A_a$, the $SO(3)$ generators in the presence of t'Hooft-Polyakov monopole can be written as²⁶

$$L_{ab} = \Lambda_{ab} - id^2 f_{ab}, \quad a = 1, 2, 3 \quad (12)$$

where $\Lambda_{ab} = -id_a D_b + id_b D_a$ and $f_{ab} = -iF_{ab}^c \frac{\sigma_c}{2}$. Defining $I_a = \frac{1}{2} \epsilon_{abc} L_{bc}$, one finds easily that $[I_a, I_b] = i\epsilon_{abc} I_c$ satisfying the $SO(3)$ algebra. Using Eq.(8) and Eq.(9), one can show

$$L_{ab} = L_{ab}^{(0)} + \epsilon_{abc} \frac{\sigma_c}{2} \quad (13)$$

where $L^{(0)}$ is the orbital angular momentum, defined by $L_{ab}^{(0)} = -id_a\partial_b + id_b\partial_a$. Define²⁴

$$V = \exp \left[i \frac{\vartheta \sigma_{3a} d_a}{\sqrt{d^2 - d_3^2}} \right] \quad (14)$$

where $\sigma_{ab} = \epsilon_{abc}\sigma_c/2$ and $\cos \vartheta = d_3/d$. One can perform a singular $SO(3)$ gauge transformation on L_{ab} such that $J_{ab} = VL_{ab}V^\dagger$ where

$$J_{\mu\nu} = -id_\mu\partial_\nu + id_\nu\partial_\mu + \epsilon_{\mu\nu}\frac{\sigma_3}{2} \quad (15)$$

$$J_{\mu 3} = -id_\mu\partial_3 + id_3\partial_\mu - \epsilon_{\mu\nu}\frac{d_\nu}{d+d_3}\frac{\sigma_3}{2} \quad (16)$$

with $\mu, \nu = 1, 2$. $\epsilon_{\mu\nu}$ is the antisymmetry tensor which has only one component $\epsilon_{12} = 1$ in this case. It is obvious that $J_{\mu\nu} = J_{12}$ forms the $U(1)$ generator on S^2/Z_2 . From the definition of Eq.(12), one can extract the $U(1)$ BPC regardless the unnecessary $\frac{\sigma_3}{2}$.

$$a_\mu = -i\epsilon_{\mu\nu}\frac{d_\nu}{d(d+d_3)}, \quad a_3 = 0 \quad (17)$$

and the finite $U(1)$ field strength over S^2/Z_2

$$F_{ab} = i\epsilon_{abc}\frac{d_c}{d^3} \quad (18)$$

We should notice that the singular gauge transformation we used has a singularity string along the negative z -axis. Therefore, while the covariant $SO(3)$ BPC is finite over the whole $\hat{d}$ field, the $U(1)$ BPC has a singularity string which is reflected through the transformation. This transformation is only valid on the northern hemisphere including the equator of the S^2/Z_2 . One is able to choose another gauge which has the singularity along the positive z -axis to describe the transformation on the southern hemisphere.

The role of this singular gauge transformation is very intriguing. We can view the covariant $SO(3)$ gauge potential A_c^a in Eq.(8) as a vector d_c pointing in the isospin space. The singular gauge transformation is nothing but the rotation of the spin vector from d_c to d_3 . Therefore, the invariant subgroup of $SO(3)$ is emerged from the isometry group of the equator of S^2/Z_2 , which is $U(1)$. This mechanism accounts for the appearance of the $U(1)$ Berry phase in this problem. As a result, the $SO(3)$ Berry phase in this system is essentially equivalent to the $U(1)$ Berry phase.

III. s -WAVE QUINTET PAIRING CONDENSATE IN SPIN-3/2 SYSTEM

A. Goldstein manifold and the 2^{nd} Hopf map

Another candidate for non-trivial gauge structures is the spin-3/2 fermionic system with contact interaction, in which an exact $SO(5)$ symmetry was identified

recently¹⁸. It may be studied in the ultra-cold atomic systems, like ^9Be , ^{132}Cs , ^{135}Ba , ^{137}Ba . The four-component spinor $(c_{\frac{3}{2}}, c_{\frac{1}{2}}, c_{-\frac{1}{2}}, c_{-\frac{3}{2}})^T$ forms the spinor representations of the $SU(4)$ group which is the unitary transformation of the four-component complex spinor. The kinetic energy term has the explicit $SU(4)$ symmetry. However, the s -wave contact interaction term breaks the $SU(4)$ symmetry to $SO(5)$. Because of the s -wave scattering, there are only the singlet and quintet channels as required by the Pauli's exclusion principle. Interestingly, the spin $SU(2)$ singlet and quintet channels interaction can also be interpreted as $SO(5)$ group's singlet and 5-vector representations.

The Cooper pair structures has also been studied in spin-3/2 system^{18,27}. The singlet and quintet pairing channel operators are described by

$$\begin{aligned} \eta^\dagger(r) &= \frac{1}{2}c_\alpha^\dagger(r)R_{\alpha\beta}c_\beta^\dagger(r) \\ \chi_a^\dagger(r) &= \frac{-i}{2}c_\alpha^\dagger(r)(\Gamma^a R)_{\alpha\beta}c_\beta^\dagger(r). \end{aligned}$$

where Γ^a are the $SO(5)$ Gamma matrices which takes the form

$$\Gamma^1 = \begin{pmatrix} 0 & iI \\ -iI & 0 \end{pmatrix}, \quad \Gamma^i = \begin{pmatrix} \sigma_i & 0 \\ 0 & -\sigma_i \end{pmatrix}, \quad \Gamma^5 = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix} \quad (19)$$

satisfying the Clifford algebra $\{\Gamma^a, \Gamma^b\} = 2\delta^{ab}$. The $SO(5)$ charge conjugate matrix R is given by

$$R = -i \begin{pmatrix} \sigma_2 & 0 \\ 0 & \sigma_2 \end{pmatrix} \quad (20)$$

The quintet pairing structure is spanned by the five polar like operators $\chi_{1\sim 5}^\dagger$, whose expectation value has a 5-vector and a phase structure as $d_a e^{i\phi}$. The Goldstone manifold for the quintet pairing is

$$\begin{aligned} R_{3/2} &= \frac{SO(5)_s \otimes SO(3)_L \otimes U(1)}{SO(4)_s \otimes SO(3)_L \otimes Z_2} \\ &= S^4 \otimes U(1)/Z_2, \end{aligned} \quad (21)$$

where the Z_2 symmetry comes from the combined operations $d_a \rightarrow -d_a, \phi \rightarrow \phi + \pi$.

Because the 4-component spinor forms the 7-dimensional sphere, similar to the spin 1/2 case, the adiabatic transportation of the quasi-particle around a half-quantum vortex in our spin-3/2 system defines a map

$$S^7 \longrightarrow S^4/Z_2, \quad (22)$$

which is homotopically equivalent to the second Hopf map *i.e.* $S^7 \rightarrow S^4$.

B. Berry connection, 2^{nd} Chern number, $SO(4)$ monopole and Yang monopole

Let us introduce the spinor

$$\Psi_k^\dagger = \left(c_{k, \frac{3}{2}}^\dagger, c_{k, \frac{1}{2}}^\dagger, c_{k, \frac{-1}{2}}^\dagger, c_{k, \frac{-3}{2}}^\dagger, c_{-k, \frac{3}{2}}, c_{-k, \frac{1}{2}}, c_{-k, \frac{-1}{2}}, c_{-k, \frac{-3}{2}} \right), \quad (23)$$

where $c_{k\sigma}^\dagger$ is the creation operator of an electron with the spin component σ and momentum k . The mean field Hamiltonian can be written as

$$\mathcal{H} = \sum_k \Psi_k^\dagger H_k \Psi_k \quad (24)$$

with

$$H_k = \begin{pmatrix} \epsilon_k \Gamma^0 & \Delta_k \\ \Delta_k^\dagger & -\epsilon_k \Gamma^0 \end{pmatrix} \quad (25)$$

where ϵ_k is the kinetic energy on the lattices referenced from the Fermi surface. $\Delta_k = -\Delta_k d_a \Gamma^a R$ while Δ_k contains the magnitude and the phase of the superconducting (SC) order parameter. The momentum k is summed only over half of the Brillouin zone to avoid the double counting. The subscript a runs from 1 to 5. d_a forms a 4-dimensional sphere S^4 . Γ^0 is the 4×4 identity matrix, and Γ^a are given by Eq.(19). The eigenvalues of Eq.(25) are

$$\lambda = \pm E_k = \pm \sqrt{\epsilon_k^2 + |\Delta_k|^2} \quad (26)$$

and their corresponding engenvectors are

$$\psi_\alpha^+(k) = \frac{1}{\sqrt{(E_k + \epsilon_k)^2 + |\Delta|^2}} \begin{pmatrix} (E_k + \epsilon_k)|\alpha\rangle \\ \Delta_k^\dagger |\alpha\rangle \end{pmatrix} \quad (27)$$

$$\psi_\alpha^-(k) = \frac{1}{\sqrt{(E_k + \epsilon_k)^2 + |\Delta|^2}} \begin{pmatrix} \Delta_k |\alpha\rangle \\ (E_k + \epsilon_k)|\alpha\rangle \end{pmatrix} \quad (28)$$

where $|\alpha\rangle$ are $SU(4)$ spinors.

We are interested in the system with the presence of half-quantum vortices. The formation of this kind of vortices is very similar to the ones in the spin-1/2 system. If we transport the spin-3/2 fermion adiabatically around one of them, a nontrivial phase is accumulated due to the $\hat{d}$ field. The BPC and the covariant field strength can be obtained respectively by

$$A_a = \frac{i}{d^2} d_c \Gamma^{ca}, \quad a = 1, 2, 3, 4, 5 \quad (29)$$

and

$$F_{abc} = -\frac{i}{d^3} (d_a \Gamma^{bc} + d_b \Gamma^{ca} + d_c \Gamma^{ab}) \quad (30)$$

where $\Gamma^{ab} = \frac{1}{4i} [\Gamma^a, \Gamma^b]$ making up of the $SO(5)$ generators. Similar non-abelian gauge structures also appeared

in the pseudoparticle field in high dimensions^{28,29,30}. Instead of the first Chern number, we have the non-vanishing second Chern number.

$$\mathcal{C}_2 = -\frac{1}{96\pi^2} \oint d\Omega_d \text{Tr} (F_{abc} F_{abc}) = 1. \quad (31)$$

where $d\Omega_d$ denotes the integration over the angular part of d_a . The field strength on S^4 , $f_{ab} = [D_a, D_b] = \partial_a A_b - \partial_b A_a + [A_a, A_b]$, can be obtained

$$f_{ab} = -i \frac{1}{d^2} P_{ab,cd} \Gamma^{cd} \quad (32)$$

where $P_{ab,cd} = \frac{1}{2} (\delta_{ac} \delta_{bd} - \delta_{ad} \delta_{bc} + \delta_{ad} d_b d_c - \delta_{bd} d_a d_c - \delta_{ac} d_b d_d + \delta_{bc} d_a d_d)$. $P_{ab,cd}$ is a 10×10 matrix because a and b are anti-symmetric as well as c and d . Similar to the projection operator $\delta_{\mu\nu} - \hat{q}_\mu \hat{q}_\nu$ in QED, $P_{ab,cd}$ is the transverse projection operator from 10-dimensional space to 6-dimensional space satisfying $d_a P_{ab,cd} = 0$. The relation between F_{abc} and ordinary field strength f_{ab} can be revealed if we define G_{ab} which is dual to F_{abc} by $G_{ab} = \epsilon_{abcde} F_{abc}$ ^{29,30}. Then, one can show

$$f_{ab} = \frac{1}{2} \epsilon_{abcde} \frac{d_c}{d} G_{de}. \quad (33)$$

Because of the projection operator $P_{ab,cd}$ in f_{ab} , the fundamental degrees of freedom of the gauge structure in this problem is not $SO(5)$ but $SO(4)$, because $SO(4)$ has six generators. We shall show that it is able to make a route from f_{ab} to the $SO(4)$ gauge field strength using a singular gauge transformation. Similar to the analysis in the second section, the transformation operator in this $SO(5)$ case has the following form

$$U = \exp \left[i \frac{\vartheta \Gamma^{5\mu} d_\mu}{\sqrt{d^2 - d_5^2}} \right], \quad \mu = 1, 2, 3, 4 \quad (34)$$

where $\cos \vartheta = d_5/d$. Then, Eq.(29) becomes

$$a_\mu = \frac{-i}{d(d + d_5)} \Sigma^{\mu\nu} d_\nu, \quad \mu = 1, 2, 3, 4 \quad (35)$$

$$a_5 = 0, \quad (36)$$

where $\Sigma_{\mu\nu}$ are the $SO(4)$ generators in the $(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$ representation which have the following form

$$\Sigma_{\mu\nu} = \begin{pmatrix} \eta_{\mu\nu}^i \frac{\sigma_i}{2} & 0 \\ 0 & \bar{\eta}_{\mu\nu}^i \frac{\sigma_i}{2} \end{pmatrix} \quad (37)$$

where $\eta_{\mu\nu}^i = \epsilon_{i\mu\nu 4} + \delta_{i\mu} \delta_{4\nu} - \delta_{i\nu} \delta_{4\mu}$ is the t'Hooft symbol, μ and ν runs from 1 to 4. In this reducible representation

of $SO(4)$ gauge group, one can easily distillate the $SU(2)$ ingredients because $SO(4) = SU(2) \otimes SU(2)$. The self-dual $SU(2)$ gauge field is given by

$$a_\mu = \frac{-i}{d(d+d_5)} \eta_{\mu\nu}^i d_\nu \frac{\sigma_i}{2} \quad \mu = 1, 2, 3, 4$$

$$a_5 = 0. \quad (38)$$

Similar to the spin-1/2 system, we obtain the $SO(4)$ BPC which is only defined on the northern hemisphere with the equator. The singularity string along the negative z -axis inherits from the singular gauge transformation. The role of the singular gauge transformation by U can be also interpreted as the rotation of a 5-dimensional vector from an arbitrary direction d_a to d_5 in the 5-dimensional isospin space. Therefore, the invariant subgroup is the isometry group of the equator S^3/Z_3 , which is $SO(4)$. Surprisingly, the representation we achieve in $SO(4)$ gauge theory is the reducible $(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$, which is the direct sum of two $SU(2)$ gauge theory. Thus, the $SU(2)$ Berry phase naturally arises in this system.

This $SU(2)$ nature of the Berry phase in the spin-3/2 system is manifested if we choose a special spinor $|\alpha\rangle$ such that $\Delta_{\mathbf{k}}|\alpha\rangle = |\Delta_{\mathbf{k}}|e^{-i\phi_\alpha}|\alpha\rangle$, which is studied by Demler and Zhang¹⁴. In this representation, $|\alpha\rangle$ is not only a $SU(4)$ spinors but also a $SO(5)$ one. Then, the BPC is given by

$$A_i^{\pm\alpha\beta} = \langle\psi_\alpha^\pm|\partial_i|\psi_\beta^\pm\rangle = \langle\alpha|\partial_i|\beta\rangle. \quad (39)$$

This is exactly the $SU(2)$ holonomy in the context of Demler and Zhang¹⁴. The special choice of the spinors $|\alpha\rangle$ is equivalent to fixing $\hat{d}_a = \hat{d}_5$ in our notion.

IV. CONCLUSION AND DISCUSSION

In summary, we found that the $SU(2)$ non-Abelian Berry phase emerges naturally in quintet condensates of

spin 3/2 fermions. The underlying algebraic structures for the ${}^3\text{He}$ and the spin-3/2 system are the 1st and the 2nd Hopf map respectively. The Chern numbers for both cases were obtained in a standard manner. In the previous case, only the first Chern number is non-vanishing, while in the later case, only second Chern number is non-zero. Both systems appear to have *finite* gauge potential, which means that the BPC can be defined covariantly everywhere over the d -field. However, the corresponding $U(1)$ and $SU(2)$ gauge connections can only be defined patch by patch in the d -space. The bridge across between the finite gauge connection and the one with singularity is constructed by the singular gauge transformation, which can only be defined patch by patch in the d -space as well³¹. The singular gauge transformation also bear with some physical meaning. It can be understood as the rotation in the spin (isospin) space, namely d_a . When the spin (isospin) points to the north pole, the invariant subgroup becomes the isometry group of the equator. When it is rotated by the gauge transformation, the gauge structure becomes finite and covariant over the whole d space.

To experimentally manifest this topological effect, the spin-3/2 ultra-cold atomic systems may serve as a promising candidate. It may also shed some light on measuring the second Chern number, which has not been revealed by any system so far. Furthermore, our calculation can also be generalized to consider the spin- $\frac{7}{2}$ superconductors in which the algebraic structure is suggested to be the third Hopf map³².

This work is supported by the NSF under grant numbers DMR-0342832 and the US Department of Energy, Office of Basic Energy Sciences under contract DE-AC03-76SF00515. HDC and CJW are also supported by the Stanford Graduate Fellowship.

-
- ¹ M. V. Berry, Proc. R. Soc. London A **392**, 45 (1984).
 - ² D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. Denny, Phys. Rev. Lett. **49**, 405 (1982).
 - ³ M. Kohmoto, Annals of Physics **160**, 343 (1985).
 - ⁴ S. C. Zhang, T. H. Hansson, and S. Kivelson, Phys. Rev. Lett. **62**, 82 (1989).
 - ⁵ A. Auerbach, *Interacting Electrons and Quantum Magnetism* (Springer-Verlag, New York, 1994).
 - ⁶ F. D. M. Haldane, Phys. Rev. Lett. **50**, 1153 (1983).
 - ⁷ F. D. M. Haldane, Phys. Lett. A **93**, 464 (1983).
 - ⁸ G. Volovik, *Exotic properties of superfluid ${}^3\text{He}$* (World Scientific, Singapore, New Jersey, London, Hong Kong, 1992).
 - ⁹ N. Read and D. Green, Phys. Rev. B **61**, 10267 (2000).
 - ¹⁰ Y. Hatsugai, S. Ryu, and M. Kohmoto, cond-mat/0308332.
 - ¹¹ F. Wilczek and A. Zee, Phys. Rev. Lett. **52**, 2111 (1984).
 - ¹² A. Zee, Phys. Rev. A **38**, 1 (1988).
 - ¹³ J. E. Avron, L. Sadun, J. Segert, and B. Simon, Phys. Rev. Lett. **61**, 1329 (1988).
 - ¹⁴ E. Demler and S. C. Zhang, Annals of Physics **271**, 83 (1999).
 - ¹⁵ S. C. Zhang and J. P. Hu, Science **294**, 823 (2001).
 - ¹⁶ S. Murakami, N. Nagaosa, and S. C. Zhang, Science **301**, 1348 (2003).
 - ¹⁷ S. Murakami, N. Nagaosa, and S.-C. Zhang, cond-mat/0310005.
 - ¹⁸ C. J. Wu, J. P. Hu, and S. C. Zhang, Phys. Rev. Lett. **91** (2003).
 - ¹⁹ P. Hilton, *An Introduction to Homotopy Theory* (Cambridge Univ. Press, Cambridge, England, 1953).
 - ²⁰ J. B. Yau, E. P. D. Poortere, and M. Shayegan, Phys. Rev. Lett. **88** (2002).
 - ²¹ M. Yang, C. Yang, K. Cheng, and Y. Lyanda-Geller, cond-mat/0208260.
 - ²² S. Weinberg, *The Quantum Theory of Fields: Modern*

- Applications (Volume II)* (Cambridge Univ. Press, Cambridge, England, 1996).
- ²³ D. G. Boulware, L. S. Brown, R. N. Cahn, S. D. Ellis, and C. Lee, Phys. Rev. D **14**, 2708 (1976).
 - ²⁴ Y. Ohnuki and S. Kitakado, J. Math. Phys. **34**, 2827 (1993).
 - ²⁵ C.-H. Chern, J.-P. Hu, and S.-C. Zhang, unpublished.
 - ²⁶ C. N. Yang, J. Math. Phys. **19**, 2622 (1978).
 - ²⁷ T. L. Ho and S. Yip, Phys. Rev. Lett. **82**, 247 (1999).
 - ²⁸ R. Jackiw and C. Rebbi, Phys. Rev. D **14**, 517 (1976).
 - ²⁹ S. L. Adler, Phys. Rev. D **6**, 3445 (1972).
 - ³⁰ S. L. Adler, Phys. Rev. D **8**, 2400 (1973).
 - ³¹ J. Moody, A. Shapere, and F. Wilczek, Phys. Rev. Lett. **56**, 893 (1986).
 - ³² B. A. Bernevig, J. P. Hu, N. Toumbas, and S. C. Zhang, Phys. Rev. Lett. **91** (2003).